

REVIEW



Pseudomonad reverse carbon catabolite repression, interspecies metabolite exchange, and consortial division of labor

Heejoon Park^{1,3} · S. Lee McGill^{2,3} · Adrienne D. Arnold^{2,3} · Ross P. Carlson^{1,2,3} 

Received: 4 November 2019 / Revised: 4 November 2019 / Accepted: 12 November 2019 / Published online: 25 November 2019
© Springer Nature Switzerland AG 2019

Abstract

Microorganisms acquire energy and nutrients from dynamic environments, where substrates vary in both type and abundance. The regulatory system responsible for prioritizing preferred substrates is known as carbon catabolite repression (CCR). Two broad classes of CCR have been documented in the literature. The best described CCR strategy, referred to here as classic CCR (cCCR), has been experimentally and theoretically studied using model organisms such as *Escherichia coli*. cCCR phenotypes are often used to generalize universal strategies for fitness, sometimes incorrectly. For instance, extremely competitive microorganisms, such as Pseudomonads, which arguably have broader global distributions than *E. coli*, have achieved their success using metabolic strategies that are nearly opposite of cCCR. These organisms utilize a CCR strategy termed ‘reverse CCR’ (rCCR), because the order of preferred substrates is nearly reverse that of cCCR. rCCR phenotypes prefer organic acids over glucose, may or may not select preferred substrates to optimize growth rates, and do not allocate intracellular resources in a manner that produces an overflow metabolism. cCCR and rCCR have traditionally been interpreted from the perspective of monocultures, even though most microorganisms live in consortia. Here, we review the basic tenets of the two CCR strategies and consider these phenotypes from the perspective of resource acquisition in consortia, a scenario that surely influenced the evolution of cCCR and rCCR. For instance, cCCR and rCCR metabolism are near mirror images of each other; when considered from a consortium basis, the complementary properties of the two strategies can mitigate direct competition for energy and nutrients and instead establish cooperative division of labor.

Keywords Diauxie · Carbon catabolic repression · Reverse carbon catabolic repression · Division of labor · Overflow metabolism

Introduction

Most natural environments are physically, chemically, and temporally complex, with the resident microorganisms exposed to multifactorial selection pressures. It is not possible to optimize all cellular functions simultaneously due to constraints on cellular resources such as anabolic nitrogen or cytoplasmic volume; a concept illustrated by the ‘Darwinian Demon’ thought exercise [1, 2]. Therefore, microorganisms

must dynamically change their phenotypes with environmental fluctuations to maintain fitness [3]. Metabolic regulation of substrate-consumption order is paramount to fitness. There are a few broad strategies that can be utilized when an organism is presented with two potential substrates: (1) utilize the ‘optimal’ substrate, (2) utilize the less ‘optimal’ substrate, or (3) co-catabolize both substrates simultaneously (see Box 1 for discussion of ‘optimal’ substrates). It can be more resource effective to catabolize a single substrate at a time as opposed to expressing multiple catabolic pathways simultaneously [4, 5]; although, exceptions exist [6]. Carbon catabolite repression (CCR) is a global regulation system that controls the sequential catabolism of preferred substrates from a milieu (Box 2). CCR is associated with generalist microorganisms that can utilize multiple substrates and exist in environments, where these substrates are often available at varying abundances

✉ Ross P. Carlson
rossc@montana.edu

¹ Department of Chemical and Biological Engineering,
Montana State University, Bozeman, USA

² Department of Microbiology and Immunology, Montana
State University, Bozeman, USA

³ Center for Biofilm Engineering, Montana State University,
Bozeman, USA

Box 1: 'Optimal' substrates.

Monod's study of *E. coli* as the model organism has influenced what is generally considered 'optimal behavior' in computational biology. Species expressing cCCR have formed the basis of much of modern systems biology studies and their metabolic regulation is the basis of textbooks [26]. Glucose is often considered the optimal substrate preferred by chemoheterotrophs, but the notion of an optimal substrate is worthy of further discussion because no substrate is universally preferred by all microorganisms. The monosaccharide has competitive properties including the flexible production of cellular energy from fermentation, respiration or combinations of the two. Glucose integrates, with nominal enzymatic steps, into central metabolism pathways including the Embden-Meyerhof-Parnas (EMP), Entner-Doudoroff (ED) and pentose-phosphate (PP) pathways. Glucose provides convenient ratios of reducing equivalents to metabolic intermediates necessary for biosynthetic fluxes. However, oligomers like cellobiose and lactose are preferred to glucose by some fermenting anaerobes. Fermenters are often considered cellular energy poor and the transport cost of substrate can be a substantial energy drain [27]. The disaccharides are more energetic on a per molecule basis than glucose, so using an ABC transporter to import a disaccharide has a better energetic payback than using an ABC transporter to import glucose [28, 29]. This strategy comes with a tradeoff, as the cell requires investment into additional enzymes to hydrolyze, phosphorylate, and in the case of lactose, process the intermediates so they can enter glycolysis. rCCR organisms like *P. putida* and *P. aeruginosa* prefer citrate, succinate, lactate, and acetate over glucose even though these substrates contain less energy per molecule and are not generally fermentable [9]. The necessary respiratory metabolism requires large investments into tricarboxylic acid (TCA) cycle and electron transport chain enzymes as well as a supply of terminal electron acceptor like O₂. Citrate and succinate are more oxidized than glucose which would reduce the electron flux per carbon mole of substrate oxidized. The *Pseudomonas* growth rate on the organic acids is not necessarily faster than on glucose [30].

These non-glucose substrates are certainly 'optimal' substrates if the appropriate ecological criteria are considered. The designation of glucose as the 'optimal' substrate is analogous to outdated arguments describing an overflow metabolism as a 'wasteful metabolism'; in recent years, this metabolic phenotype has been appreciated as a competitive ecological strategy [31, 32].

Preferred substrate	Formula	ΔG combustion (kJ mol ⁻¹)*	Degree of reduction	Example microorganism	Strategy
Cellobiose	C ₁₂ H ₂₂ O ₁₁	-5652 (s)	4	<i>C. thermocellum</i>	Cost effective transport, fermentable
Lactose	C ₁₂ H ₂₂ O ₁₁	-5652 (s)	4	<i>B. longum</i>	Cost effective transport, reciprocal substrate, fermentable
Glucose	C ₆ H ₁₂ O ₆	-2805 (s)	4	<i>E. coli</i> , <i>B. subtilis</i>	Cost effective resource investment
Citrate	C ₆ H ₈ O ₇	-1920 (s)	3	<i>P. putida</i> , <i>P. aeruginosa</i>	Reciprocal substrate? Division of labor?
Succinate	C ₄ H ₆ O ₄	-1491 (c)	3.5	<i>P. putida</i> , <i>P. aeruginosa</i>	Reciprocal substrate? Division of labor?
Lactate	C ₃ H ₆ O ₃	-1368 (l)	4	<i>P. putida</i> , <i>P. aeruginosa</i>	Reciprocal substrate? Division of labor?
Acetate	C ₂ H ₄ O ₂	-874 (l)	4	<i>P. putida</i> , <i>P. aeruginosa</i>	Reciprocal substrate? Division of labor?

*Combustion values from Bioprocess Engineering Principles, Doran, 2013 ed 2. Academic Press. Combustion value for cellobiose assumed equivalent to lactose.

The cCCR regulation scheme, known under specific conditions as diauxie, has been the focus of much experimental and theoretical research [7–12]. The historical basis of cCCR goes back more than 70 years to Jacques Monod and his observations of *Escherichia coli* growing in the presence of glucose and other sugars [13]. Monod recorded *E. coli* cultures growing exponentially on glucose until depleted,

followed by a lag phase, then a second exponential growth phase on the other sugar. Diauxic growth is a trade-off between specializing for a single nutrient and being able to switch between nutrients as conditions change [14]. Committing to one metabolism at a time, necessitates a metabolic shift when the preferred substrate is exhausted, with a resulting temporal delay.

Box 2: Carbon catabolite repression.

Carbon catabolite repression (CCR): A regulatory network which selects ‘preferred’ carbon sources from a pool of potential carbon sources. The ecological basis of a preferred carbon source is environment dependent and Nature has evolved numerous strategies for optimizing the catabolism of available carbon sources. The best studied and often default example of CCR is that of *E. coli* or *B. subtilis*, which prefer sugars like glucose to other potential carbon sources like organic acids. These model organisms have strongly influenced the criteria applied to identify ‘optimal’ phenotypes in computational biology studies of chemoheterotrophs. However, as science describes a broader range of microbiological physiologies from diverse environments, the number of different CCR strategies is sure to expand. In this review, we consider primarily two CCR strategies defined below.

Classic carbon catabolite repression (cCCR): The textbook example of CCR, first described in model organism *E. coli*, is called cCCR here to distinguish it from the general term CCR and other specific CCR strategies. cCCR phenotypes prefer glucose to other carbon substrates [26].

Reverse carbon catabolite repression (rCCR): A CCR strategy first described in *Pseudomonads* in 1959 and later termed ‘reverse’ diauxie or rCCR because the hierarchy of ‘preferred’ carbon sources was nearly ‘reverse’ that of cCCR preferences [9,33].

The traditional interpretation of diauxie as being carried out by a single population of organisms, synchronized in function, that sequentially catabolizes preferred and non-preferred substrates has been challenged [15, 16]. Recent research has posited an alternative interpretation; rather than a homogenous population, the observed lag phase can be explained by dynamic shifts in subpopulations possessing different fitness. For instance, during growth on glucose, multiple subpopulations catabolize the sugar but upon glucose depletion only some subpopulations, in an ecological bet-hedging strategy, are phenotypically prepared to consume the less desirable substrate [17]. The lag in growth between the two sugars corresponds to a shift in subpopulation distributions as fit subpopulations outgrow other subpopulations [17]. Control of these subpopulations has been associated with the regulation of bistable genetic switches [18, 19].

cCCR is not used by all glucose-utilizing microorganisms. *Pseudomonads* and *Acinetobacter* [20] utilize substrates in an order almost reverse of cCCR. Therefore, this metabolic regulatory strategy was termed reverse carbon catabolite repression (rCCR) or sometimes reverse diauxie. rCCR utilizing microorganisms are very competitive as evidenced by their broad global distributions and are arguably at least as competitive as the better studied members of the *Enterobacter* and *Firmicutes* used to define cCCR. The ecological rationale of the reverse diauxie strategy is still an open question and is discussed in later sections.

The study of cCCR and rCCR has traditionally focused on monocultures for the sake of simplicity, possibly limiting the interpretation of regulatory systems, as most microorganisms have existed on evolutionary timeframes within consortia [21–24]. The organisms have interacted with other species for millennia, and these interactions have influenced access to energy and nutrients [25]. Many genes regulated

by the CCR systems such as quorum sensing, biofilm formation, and antibiotic or reactive oxygen species (ROS) tolerance have been studied from the perspective of medical challenges and monocultures, even though these behaviors modulate inter-microbial interactions and strategies for resource acquisition in consortia (see sections below).

This review summarizes (1) cCCR and rCCR molecular mechanisms; (2) theory used to interpret and model CCR phenotypes; and (3) the nexus of CCR, metabolism, and cellular interactions. Collectively, the properties of cCCR and rCCR phenotypes can enhance growth and persistence by enabling cooperation via division of labor, as observed in some medical environments such as chronic wounds, where cCCR and rCCR microorganisms commonly cooccur (Fig. 1).


Molecular mechanisms of CCR

Most CCR studies have been performed in model organisms such as *E. coli* and *Bacillus subtilis*, which prefer glucose over other substrates. The regulatory outcomes of CCR in *E. coli* and *B. subtilis* are similar, though the mechanisms differ (Fig. 2). This review summarizes molecular mechanisms of CCR for context and to highlight similarities and differences between cCCR and rCCR (Table 1). This summary is not exhaustive; there are numerous excellent reviews which discuss finer aspects of CCR molecular biology [7–12].

Molecular mechanisms of cCCR

Gram-negative *E. coli* utilizes two major cCCR regulation mechanisms: (1) inducer exclusion [9, 44, 47, 48] and (2) cyclic AMP (cAMP)-catabolite repressor protein (CRP) regulation of gene promoter activity; although the condition-specific importance of each mechanism is still an

Fig. 1 Illustration of classic carbon catabolite repression (cCCR) phenotype with overflow metabolism and reverse carbon catabolite repression (rCCR) phenotype as well as an illustration of the complementary relationship between the two global metabolism regulators. cCCR microorganisms prefer glucose to other substrates, while rCCR microorganisms prefer organic acids to glucose



open research question [49, 50]. Glucose can enter the cell using the multisubunit phosphoenolpyruvate (PEP): carbohydrate phosphotransferase system (PTS). Regulation of cCCR begins with the phosphorylation state of the glucose-specific subunit of the PTS transporter, EIIA^{Glc}. EIIA^{Glc} is phosphorylated by PEP via a chain of phosphorylation reactions. In the presence of high glucose flux, the phosphate from EIIA^{Glc} is transferred to the hexose producing glucose-6-phosphate, precluding the phosphate group from activating PTS EIIA components specific for less preferred substrates. This process is known as inducer exclusion, because the excluded, non-preferred substrates are inducers for their own catabolic operons [51]. When glucose flux is low, phosphorylated EIIA^{Glc} accumulates and activates EIIA subunits for less preferred substrates and activates the enzyme adenylate cyclase [35, 52] which catalyzes the production of cAMP [53]. cAMP binds to the catabolite gene activator protein (CAP), forming the cAMP–CAP complex which binds to DNA, activating the promoters of ~200 genes for a range of cellular processes [49, 50, 54–57]. The classic example of the CCR inducer exclusion and cAMP–CRP regulation occurs during glucose–lactose diauxic growth [58–61].

CCR is complex, with many levels of modulation in addition to inducer exclusion and cAMP–CRP regulation. Small regulatory RNAs (sRNA) also contribute to post-transcriptional regulation of CCR [10, 62, 63]. sRNA Spot 42 is described as a third pillar of CCR in *E. coli* [10, 64]. Spot 42 represses target genes as opposed to the role of cAMP–CRP,

which generally activates gene targets. The Spot 42 regulon has 29 documented genes [65]. Central metabolism intermediates such as α-ketoglutarate, oxaloacetate, PEP, and pyruvate also play important roles in CCR by affecting the levels of cAMP, adding additional complexity to CCR [66, 67].

E. coli cCCR regulation ‘glitches’ can result in substrate selection that does not enable the fastest growth [68]. *E. coli* cultures growing on a mixture of glucose and lactose and a poor nitrogen source (e.g., arginine, glutamate, and proline) preferentially consume glucose over lactose, but growth on glucose is slower than on lactose due to altered cAMP concentrations. Therefore, this diauxic growth pattern has relatively slow growth during the first phase and faster growth during the second phase. The effect is attributed to laboratory conditions not reflective of habitats that influenced *E. coli* selection [68].

Gram-positive *B. subtilis* also prefers glucose and utilizes the PTS system for sugar transport, although the molecular components of CCR are different than in *E. coli*. cCCR in *B. subtilis* is mediated largely by negative regulation through a repressor protein expressed in the presence of glucose. The key cCCR components include catabolite control protein A (CcpA), phosphocarrier protein (HPr), HPr kinase (HPrK), and glycolysis intermediates fructose-1,6-bisphosphate and glucose-6-phosphate [36–38]. HPr plays two important roles. First, HPr participates in the phosphorylation cascade by transferring the phosphate group from PEP to glucose, forming glucose-6-P. Second, HPr is a regulator molecule which responds to glycolysis indicators fructose-1,6-bisphosphate

Fig. 2 Molecular components of carbon catabolite repression (CCR). **a** Major molecular components of classic carbon catabolite repression (cCCR) in Gram-negative *E. coli*. **b** Major molecular components of cCCR in Gram-positive *B. subtilis*. **c** Major molecular components of reverse carbon catabolite repression (rCCR) found in *Pseudomonads*. See main text for abbreviations and details

or glucose-6-phosphate to form a complex with CcpA. The CcpA–HPr complex, stabilized with glycolysis intermediates, can bind DNA, inhibiting expression of genes associated with non-preferred substrates [69–75]. Approximately 400 *B. subtilis* genes are influenced by CCR [76, 77].


Molecular mechanisms of rCCR

cCCR-mediated preference for glucose is not a universal strategy for generalist, glucose-catabolizing heterotrophs [9, 78, 79]. *Pseudomonas aeruginosa* and *Pseudomonas putida* consume preferred carbon substrates in nearly reverse order from *E. coli* and *B. subtilis*. They preferentially catabolize organic acids and amino acids before catabolizing glucose [8, 78, 80–82]. Study of rCCR lags the study of cCCR, highlighting a research area of opportunity.

Known rCCR mechanisms in *Pseudomonads* operate at the mRNA level; this is contrary to most cCCR microorganisms, which are governed by DNA-binding transcription regulators [83, 84]. RNA chaperone protein Hfq represses expression of catabolic targets for non-preferred substrates by binding directly to the 5' end of the mRNA. Catabolite repression control (Crc) protein interacts with Hfq, creating a stabilized Hfq/Crc/mRNA complex, enhancing regulation. Hfq and Crc regulons have significant overlap [85–87]. Computational analysis of Crc-binding sites predicted 143 candidate genes in *P. putida* and 215 candidate genes in *P. aeruginosa*, while experimental hfq gene deletion studies identified 212 target genes in *P. fluorescens* [88–90]. Small regulatory RNAs (sRNA) act as antagonists to Hfq, binding the protein and relieving repression of the non-preferred catabolic pathways. *P. aeruginosa* has 1 characterized sRNA (CrcZ), while *P. fluorescens* and *P. putida* have two (CrcY, CrcZ), as does *P. syringae* (CrcZ, CrcX) [84, 91, 92].

Theoretical aspects of CCR

CCR is the collective result of many interacting molecular components. The complexity necessitates in silico representations for quantitative predictions. Decades of in silico CCR models exist, most focusing on *E. coli* [67, 93]. The earliest in silico models were primarily mechanistic, comprised of mass balances on enzymes, metabolites, and metabolic regulators, and were represented by ordinary differential equations. As a result, they were limited by the availability of



experimentally measured enzyme parameters and metabolite concentrations [67, 93]. These models were not well-suited to address the ecological rationale of CCR strategies. One exception is cybernetic modeling, which examines implicit aspects of resource investment into metabolic strategies and can be used to study diauxic growth [95]. The modeling technique requires a priori calibration data to identify cybernetic parameters which are then used to interpret pathway resource requirements; the cybernetic parameters do not necessarily map explicitly to molecular components [96].

The omics revolution has generated huge data bases which have been leveraged by a metabolic modeling family known as stoichiometric modeling; two common types of stoichiometric modeling are flux balance analysis (FBA) and elementary flux mode analysis (EFMA) [97–100]. Stoichiometric models mathematically define the phenotypic capabilities of metabolic systems by accounting for the net transformation of all enzyme-catalyzed reactions as well as relevant abiotic reactions. The mass balances for all system components can be written as a set of linear equations when appropriate time scales are assumed; this negates the requirement for difficult to measure enzyme parameters. The modeling approach does require a minimal set of parameters such as biomass composition and cellular maintenance energy values [101].

A remarkable number of stoichiometric modeling studies have examined aspects of cCCR, including maximum growth rates, overflow metabolisms, and sequential use of substrates. Stoichiometric models define a mathematical solution space of all possible cellular phenotypes based on the linear mass balance equations; context specific flux distributions are identified using optimization criteria. The most commonly used optimization criterion identifies phenotypes that maximize growth rate, consistent with the basic tenets of cCCR [102–104]. All microbial systems are resource constrained and metabolic strategies that maximize fitness must competitively allocate limiting resources. Early applications of stoichiometric models examined limited accessibility of a catabolic resource, O_2 , and identified metabolisms that maximize energy extraction from glucose while minimizing consumption of O_2 [99, 100, 105, 106]. Resource allocation into cCCR-relevant strategies has been a topic of many stoichiometric studies and has resulted in predictions of overflow metabolisms that secrete partially oxidized byproducts such as acetate even in the presence of electron acceptors such as O_2 . Overflow metabolisms were for decades described as a wasteful metabolism that did not extract all the available energy from a substrate, but recently this view has changed [31]. Representation of resource scarcity in stoichiometric models has expanded to consider anabolic resources, and the simulations have provided ecological explanations for overflow metabolisms occurring at both fast growth rates during batch cultivation as well as slow

growth rates during chemostat cultivation [107]. Numerous proxies for anabolic cellular resources have been described in *E. coli*, with each generating similar outcomes; metabolic fluxes are directed to maximize energy production from the limiting resource, which leads to partial oxidation of substrate and an overflow metabolism. Basically, when glucose is the preferred substrate, glycolysis is an essential pathway, while auxiliary metabolic routes are optional and thus, down regulated when an anabolic resource becomes scarce. Anabolic resource considerations have included (1) accounting of the moles of a resource (e.g., nitrogen, carbon, amino acid) required to synthesize a metabolic pathway relative to the catabolic efficiency of the pathway [31, 32], (2) the volume of the cytosol occupied by enzymes relative to the catabolic efficiency of the flux distribution [103, 104], and (3) the membrane surface area occupied by membrane complexes required for a cellular flux distribution [108, 109] as well as other representations of optimal proteome investment into metabolic fluxes [5, 110]. In silico metabolism representations have also expanded to account for not only metabolic pathways but also the enzymatic machinery required to synthesize RNA and protein [111–113]. The converging of numerous resource allocation problems onto similar phenotypes highlights common relationships between the different proxies regardless of the level of molecular details considered. Resource allocation into enzymes has also been used to justify the sequential catabolizing of glucose and lactose in a mixture. For example, glucose catabolism requires fewer enzymes, permitting larger concentrations of the enzymes and driving a faster growth rate than lactose catabolism [94, 114].

The ecological basis of rCCR is not well studied. Common hypotheses invoke the availability of substrates during the organisms' evolutionary histories. In many environments, such as soils, sediments, and plant surfaces, organic acids are plentiful byproducts of biomass degradation or plant secretions [7, 8]. However, these environments also contain sugars [115]. Organisms such as *P. aeruginosa* and *P. putida* are competent glucose catabolizers, but their regulation schemes do not prioritize the substrate. A 'gluttony' hypothesis has been proposed which postulates that Pseudomonads utilize a strategy of consuming substrates as fast as possible to deny the resources to competitors [90]. The hypothesis states the rCCR regulation is necessary to balance metabolic intermediates which could accumulate to inhibitory concentrations. The theory postulates the outcome of rCCR is not to select 'preferred substrates' but rather to repress substrates in a manner that balances metabolism. The theory does not explicitly address the sequential use of substrates or why glucose is not co-metabolized.

Few, if any, modeling studies explicitly address the basic tenets of rCCR such as the preference for organic acids over glucose, the lack of almost any overflow metabolism,

and a propensity to not always maximize growth rate in monoculture [9, 30, 116–118]. Resource availability limits bacterial productivity in most environments, so widely distributed bacteria such as Pseudomonads have surely been influenced by resource scarcity. The role of resource allocation to enzymes in rCCR organisms is unknown, but it generally contradicts theories postulated for *E. coli*. rCCR organisms do not optimize investment into glycolysis, at the expense of respiration, resulting in an overflow metabolism; in fact, rCCR organisms catabolizing glucose do not typically exhibit an overflow metabolism and instead completely

oxidize the sugar to CO₂ [118–120]. All Pseudomonads catabolize glucose using the ED pathway [118, 121]. The ED pathway minimizes investment into enzymes as compared to the higher cellular-energy yielding EMP pathway [2, 31, 32, 122], (Fig. 3) providing hints of potential resource allocation strategies in Pseudomonads. In silico representations of *P. aeruginosa* and *P. putida* metabolism using stoichiometric models used the maximization of growth optimization criterion to identify flux distributions [123–126]. rCCR organisms do not always select preferred substrates based on maximizing growth rates [30].

Fig. 3 Trade-off between resource investment into enzymes and energetic efficiency of the Embden–Meyerhof–Parnas (EMP) (orange lines) or Entner–Doudoroff (ED) (blue lines) glycolysis pathways. Shared components are colored green. Pseudomonads use the ED pathway to catabolize glucose; the ED pathway requires fewer resources to assemble but also extracts less cellular energy from glucose than the EMP pathway. Reaction numbers map to enzymes, protein subunit compositions, and the resources required to assemble functional enzyme based on *E. coli* protein sequences. Data from Carlson, 2007 [31], figure modified from Carlson and Taffs, 2010 [2]




Table 1 Summary of carbon catabolite repression (CCR) mechanisms and key molecular components in representative Gram-positive and Gram-negative microorganisms

Microorganism	Gram staining	Governing mechanism	Important components	References
<i>E. coli</i>	–	Positive induction of DNA transcription	EIIA-P, cAMP, CAP	[34, 35]
<i>Bacillus subtilis</i>	+	Negative regulation of DNA transcription	CcpA, HPr, HPrK	[36–38]
<i>Lactobacillus brevis</i>	+	Negative regulation of DNA transcription	CcpA, HPr, HPrK	[39]
<i>Lactococcus lactis</i>	+	Negative regulation of DNA transcription	CcpA, HPr, HPrK	[40]
<i>Clostridium cellulyticum</i>	+	Negative regulation of DNA transcription	CcpA, HPr, HPrK	[41]
<i>Corynebacterium glutamicum</i>	+	Negative regulation of DNA transcription	Ram A, RamB	[42, 43]
<i>Acinetobacter baylyi</i>	–	Repression of mRNA translation	Crc, Hqf, CrcZ	[7–9, 44]
<i>Pseudomonas putida</i>	–	Repression of mRNA translation	Crc, Hqf, CrcZ	[7–9, 44]
<i>Pseudomonas aeruginosa</i>	–	Repression of mRNA translation	Crc, Hqf, CrcZ	[7–9, 44]
<i>Streptomyces coelicolor</i>	+	Biochemical state of glucose kinase	Glc	[45, 46]

CCR and social interactions

Microbial consortia are ubiquitous in nature, and competition for resources often predicts the outcome of microbial interactions [127–130]. Therefore, it is logical that global regulation strategies such as CCR are linked to social behaviors such as quorum sensing, biofilm formation, and virulence mechanisms including antibiotic or ROS sensitivity [131, 132]. Examples of the different social interactions for cCCR and rCCR organisms are summarized in Table 2.

Quorum sensing is an intercellular communication mechanism which enables coordination of cellular phenotypes across populations and space [133]. Quorum sensing involves the synthesis and secretion of small soluble molecules that accumulate in the environment. At critical concentrations, the soluble molecules are imported into the cells and initiate gene regulation. Quorum sensing is regulated by CCR [134–136] (Table 2). CCR-influenced quorum sensing is also required for other social behaviors including the development of mature biofilms [137, 138].

Biofilms are microbial communities encapsulated in self-produced polymer and are often associated with interfaces [139, 140]. Biofilms are social structures with high densities of cells (~300 g/L), up to two orders of magnitude higher than typical in vitro culture and many orders of magnitude higher than common environmental populations [141–143]. Metabolite transport in biofilms is primarily mediated by diffusion, and high cell density and high cellular activity create mass transfer limitation and nutritional gradients [82]. Biofilms are argued to promote the cellular interaction strategy of altruism [131, 144]. Recent research demonstrated

a remarkable substrate ‘flux time-sharing’ and metabolic oscillations used by altruistic *B. subtilis* cultures to partition resources between cells on the periphery of a biofilm and cells deep within a biofilm [145, 146]. The cells on the interior and exterior of the biofilm oscillate their activity in a coordinated manner, taking turns metabolizing the limiting substrate, which resolves conflict for the substrate and ultimately enhances community growth.

Virulence mechanisms refer to a broad range of cellular strategies that, in a host, enhance resource acquisition, growth, and persistence and are often associated with CCR [7, 147]. Common examples of virulence mechanisms include antibiotic tolerance, production of enzymes such as catalase, superoxide dismutase, and other enzymes for either producing or mitigating ROS, secretion of resource acquisition molecules such as siderophores, proteases, lipases, elastinases, or haemolysins, as well as the quorum sensing and biofilm phenotype mentioned previously (Table 2). Virulence mechanisms, such as antibiotic tolerance, are typically studied from the perspective of monocultures and medical challenges [148–150]. Antibiotics in microbial environments precede human use; these agents are part of natural microorganism interactions used to influence the distribution of species in consortia; secretion and degradation of antibiotics can stabilize consortia [151, 152]. Antibiotic and ROS susceptibility is well documented as a function of CCR [153, 154].

Table 2 Summary of social behaviors regulated by carbon catabolite repression (CCR) mechanisms in both classic (c) and reverse (r) CCR utilizing microorganisms

	Microorganism (gram stain)	CCR	Social behavior linkage to CCR	
Biofilm formation	<i>E. coli</i> (-)	c	cAMP-CRP regulates <i>csgD</i> and biofilm formation	[155, 156]
	<i>P. aeruginosa</i> (-)	r	Crc is required for aerobic biofilm formation	[157]
	<i>P. aeruginosa</i> (-)	r	CrcZ cross-regulation controls anaerobic biofilm formation	[158]
	<i>P. aeruginosa</i> (-)	r	<i>cbrB</i> is linked to biofilm formation and dispersal	[159]
	<i>S. aureus</i> (+)	c	CcpA regulates biofilm formation	[160]
	<i>S. epidermidis</i> (+)	c	CcpA regulates biofilm formation	[161]
Quorum sensing	<i>E. coli</i> (-)	c	Cocrystal structures of LsrK and HPr establishes link between CCR and QS	[162]
	<i>E. coli</i> (-)	c	cAMP-CRP influence <i>AI2</i> synthesis and uptake	[163, 164]
	<i>P. aeruginosa</i> (-)	r	Lon and CIPXP proteases link CCR and QS	[136]
	<i>S. aureus</i> (+)	c	CcpA upregulates Agr which is an autoinducing QS peptide	[165]
Stress tolerance	<i>P. aeruginosa</i> (-)	r	Hfq-dependent antibiotic susceptibility	[148]
	<i>P. aeruginosa</i> (-)	r	Oxidative stress tolerance orchestrated by Crc	[149]
	<i>E. coli</i> (-)	c	Acetate CCR regulates biofilm formation and virulence	[150]
	<i>S. aureus</i> (+)	c	CCpE control virulence factors including alpha toxin	[166]
	<i>P. syringae</i> (-)	r	Crc influences virulence, oxidative tolerance, and biofilm formation	[167]

Division of labor and resource partitioning in microbial consortia

Interactions between microorganisms with similar catabolic preferences are often antagonistic [24]. Resource partitioning through mechanisms such as division of labor or reciprocal prioritization of substrates can mitigate direct competition [22, 23, 128, 168–170]. cCCR and rCCR utilizing organisms are often found to coexist in medical environments, including chronic wounds and cystic fibrosis lungs [171]. rCCR organism *P. aeruginosa* has been isolated from 15 to 80% of chronic wounds, while cCCR organism *S. aureus* has been found in more than 90% of chronic leg ulcers [172–178]. *S. aureus* and *P. aeruginosa* are excellent biofilm formers and often isolated together [179, 180]. Synergistic interactions in polymicrobial infections can lead to greater virulence likely through enhanced resource acquisition and greater efficiency of converting resources into new biomass. For instance, wounds colonized by polymicrobial consortia have more negative patient outcomes than wounds colonized by monocultures [173, 177, 181, 182].

CCR regulates a wide range of social behaviors and likely modulates division of labor or reciprocal substrate prioritization (Table 2). cCCR and rCCR consortia represent an intuitive division of labor, whereby the rCCR organism consumes byproducts of the cCCR metabolism removing metabolites that could be a thermodynamic constraint on metabolism as well as inhibitory [183] (Fig. 1). This positive feedback mechanism would enable consortia to increase biomass productivity by more completely depleting available substrates than a monoculture [184, 185] (Fig. 3).

rCCR microorganisms prefer non-fermentable substrates such as organic acids necessitating the presence of a terminal electron acceptor, such as O_2 [116]. The availability of O_2 is a fitness challenge especially in diffusion-limited biofilms, where other organisms could consume the catabolic substrate. *P. aeruginosa* is a very competitive microorganism with mechanisms to acquire limited resources such as O_2 . First, *P. aeruginosa* utilizes quorum sensing and other community surveillance tactics to assess and regulate the type of competitive strategies employed [21, 176, 186–188]. These strategies involve the secretion of phenazine moieties such as pyocyanin, quinolones, and cyanide. The phenazine- and quinolone-based molecules have numerous social and resource acquisition activities including antibacterial properties, biofilm formation, iron chelation, cellular communication, extracellular electron transport and the production of ROS [189–193]. These molecules, along with cyanide which is also secreted by *P. aeruginosa*, can inhibit respiration in neighboring cells [188, 194]. Collectively, these mechanisms would modulate the phenotypes of nearby cells, forcing them into a fermentative metabolism which partitions O_2 for

P. aeruginosa respiration, while at the same time producing organic acids, a preferred carbon source. Long-term exposure to quinolones selects for small colony variants (SCVs) of *S. aureus* which have impaired respiratory activity and utilize primarily fermentative metabolisms [180, 188, 195].

Direct measurement of biomass productivity in natural consortia with cooccurring cCCR and rCCR microorganisms is difficult; artificial, in vitro culturing conditions often result in imbalanced resource partitioning, competition for resources and antagonism [181, 196]. Quantitative biomass productivity results are available for both evolved and synthetic systems that behave analogously to a cCCR + rCCR consortium. These systems have defined division of labor consisting of a primary, glucose-catabolizing population and a secondary, byproduct catabolizing population. The primary population prefers glucose, maximizes growth rate, and utilizes an overflow metabolism with the secretion of acetate and other byproducts. The byproduct population catabolizes the overflow byproducts and increases system productivity via two mechanisms. First, it modifies the local environment by removing acetate, which is a potent growth inhibitor that also lowers biomass yields on substrate, and secondly, the population utilizes non-preferred substrates that would otherwise be unutilized by the cCCR microorganism. The system design is simple but has a remarkable effect on system productivity. The in vitro consortium with evolved division of labor has 15% improvement in biomass productivity relative to the original monoculture during planktonic growth [197, 198]. The synthetic consortium with engineered division of labor has a 20% improvement in biomass productivity during planktonic growth and a 50% increase in biomass productivity during biofilm growth [199]. Both the evolved and engineered systems utilize substrate more effectively, driving glucose to lower concentrations [197–199]. The increases in biomass productivity can be explained by an enhanced return on carbon investment derived from the kinetic effect of division of labor, the removal of inhibitory byproducts which promotes more complete depletion of substrate, and the catabolism of non-preferred substrates that would otherwise be wasted. The section below provides a quantitative analysis of the effects.

Theoretical analysis of division of labor enhancing biomass productivity

Enzyme flux is realized by investment into both enzymes and substrate pools. There is an optimal relationship between enzyme and substrate concentrations that minimizes the total investment necessary to drive a flux [107, 200–203]. Cross feeding consortia utilizing division of labor can have a better functional return on investment based on the properties of Michaelis–Menten kinetics. An example is presented which

demonstrates a reduced total requirement for carbon with cross feeding between two specialist consortium members as compared to two generalist microorganisms. The aggregate specialist consortium and generalist system perform the same net transformation of glucose to CO_2 and both systems consume glucose at the same total rate. For the quantitative example, enzymes from glycolysis are represented by properties of the Pgi enzyme and enzymes from the tricarboxylic acid (TCA) cycle are represented by the properties of the FumA enzyme. Irreversible Michaelis–Menten kinetics are assumed for the single substrate reactions; enzyme amino acid sequence and subunit composition for *E. coli* are used to calculate carbon investment into enzymes and substrates. Figure 4 details the enzyme parameters, overall biochemical pathways, and the cross feeding scheme based on a cCCR and rCCR metabolism.

The generalist system is comprised of two cells each operating a complete oxidation of glucose and each consuming glucose at a specific rate of $1 \text{ mmol glucose g cdw}^{-1} \text{ h}^{-1}$. The generalist cells have 21 considered enzymatic steps. The specialist consortium is also comprised of two cells with the first cell consuming glucose at a rate of $2 \text{ mmol glucose g cdw}^{-1} \text{ h}^{-1}$ and secreting lactate at $4 \text{ mmol lactate g cdw}^{-1} \text{ h}^{-1}$. The first cell has 10 considered enzymatic reactions. The second cell consumes the lactate at the same rate it is secreted. The second cell has 13 considered enzymatic reactions. When carbon investment into both enzymes and substrate pools is considered, the cross feeding system requires ~4% smaller investment to operate the same net transformation even though it contains two more enzyme-catalyzed steps than the generalist system. This system is analogous to cross feeding between *S. aureus* and *P. aeruginosa*. The reduced resource requirement is available for enhanced growth, providing an explanation for the enhanced biomass productivity.

The catabolism of overflow byproducts removes inhibitors, which improves overall consortium productivity, as

mentioned above. This effect has been quantified in a recent stoichiometric modeling study that examined biofilm growth of an engineered, division of labor consortium which cross feeds acetate [204]. The model incorporated the genome-encoded metabolic potential of the two microorganisms, the diffusion-based transport of metabolites in the biofilm, and considered the inhibitory properties of the exchanged acetate (Fig. 5). The simulations captured the enhanced biomass productivity, approximately 50% improvement, observed in the corresponding experimental study based on the removal of the inhibitory acetate [199]. This modeling study did not account for resource investment into intracellular enzymes and metabolites highlighting how each mechanism in isolation can enhance productivity. Integrating both mechanisms into a computational representation is anticipated to further capture in situ mechanisms of productivity.

Conclusions

Carbon catabolite repression is the result of natural selection within dynamic environments. Two common forms of carbon catabolite repression, classic carbon catabolite repression and reverse carbon catabolite repression have broad global distributions in natural microbial populations and have near mirror-image preferences for substrates. These metabolic strategies are typically studied from the perspective of monocultures. Here we present a consortia-based argument for the two metabolic strategies based on reciprocal social interactions including quorum sensing, biofilm formation, and virulence mechanisms. Appropriate ecological interpretation of global regulation schemes can open many new avenues of research leading to a greater understanding of the microbial world and its uses in agriculture, industry, and medicine. Box 3 highlights interesting, unresolved questions regarding rCCR

Box 3: Unresolved CCR questions.

- 1) What ecological strategy explains the intracellular resource allocation patterns observed during a rCCR?
- 2) rCCR strategies are respiration-centric so how does the phenotype allocate cellular membrane surface area relative to a cCCR microorganism utilizing an overflow metabolism?
- 3) How does the distribution of cellular membrane surface area allocated to enzymes change during division of labor and cross feeding?
- 4) *P. aeruginosa* is a proficient biofilm former and operates a respiration-centric rCCR metabolism which sets up an important ecological question, does the *P. aeruginosa* biofilm phenotype also employ a rCCR strategy even though biofilms are limited by O_2 diffusion?
- 5) How do distributions of cCCR and rCCR microorganisms change with time in chronic wounds and CF lungs? Does early colonization favor a cCCR metabolism while evolved communities select for interacting consortia?

Fig. 4 Illustration of enhanced functional return on carbon investment in a cross feeding consortium relative to a generalist system. Enzyme flux requires investment into both enzyme and metabolite pools which can be optimized to reduce total cellular investment. Operating enzymes at higher fluxes represents a better functional return on resource investment (e.g., aggregate carbon atoms in enzyme and substrate). Analysis considers Michaelis–Menten-type kinetics, 2 cells using a generalist strategy, or 2 specialist cells cross feeding. The overall glucose flux and glucose transformation are the same for both scenarios. Glycolysis reactions are represented by the enzyme Pgi, while tricarboxylic acid (TCA) cycle enzymes are represented by the enzyme FumA. Enzyme values are from *E. coli* and obtained from Brenda and EcoCyc. The specialist consortium requires a smaller investment of carbon to attain the same flux and transformation as the generalist system. Specific glucose uptake rate (q_1) was set to 1 mmol glucose g cdw⁻¹ h⁻¹. Portion of figure modified from Beck et al. 2016 [200]





Fig. 5 Role of byproduct inhibition in cross feeding consortia growing as biofilms. **a** Schematic of metabolic interactions between cCCR- and rCCR-cell types. **b** Example of the inhibitory properties of an organic acid as a function of concentration. **c** *In vitro* data for an engineered, cross feeding consortium growing as a biofilm. **d** *In silico* prediction of biomass productivity for an engineered consortium growing as a biofilm. **e** Micrograph of a cryosectioned biofilm comprised of an engineered consortium, rCCR-cell type expressing rfp, cCCR-cell type expressing gfp. **f** *In silico* predictions of cell-type spatial distributions within a biofilm. Portions of the figure are modified from Patel et al. 2019 [204] and Bernstein et al. 2012 [199]



Acknowledgements HP was supported by the Army Research Office award W911NF-16-1-0463, SLM was supported by the National Institutes of Health award U01EB019416, ADA was supported by the National Science Foundation award OIA 1736255, and RPC was supported by National Science Foundation award 1361240.

References

- Law R (1979) Optimal life histories under age-specific predation. *Am Nat* 114(3):399–417. <https://doi.org/10.1086/283488>
- Carlson RP, Taffs RL (2010) Molecular-level tradeoffs and metabolic adaptation to simultaneous stressors. *Curr Opin Biotechnol* 21(5):670–676
- Shoval AO, Sheftel H, Shinar G, Hart Y, Ramote O, Mayo A, Dekel E, Alon U, Science S, Series N, June N (2012) Evolutionary trade-offs, pareto optimality, and the geometry of phenotype space. *Science* 336(6085):1157–1160
- Sandberg TE, Lloyd CJ, Palsson BO, Feist AM (2017) Laboratory evolution to alternating substrate environments yields distinct phenotypic and genetic adaptive strategies. *Appl Environ Microbiol* 83(13):e00410–e00417. <https://doi.org/10.1128/AEM.00410-17>

5. Wang X, Xia K, Yang X, Tang C (2019) Growth strategy of microbes on mixed carbon sources. *Nat Commun* 10(1):1279. <https://doi.org/10.1038/s41467-019-09261-3>
6. Frunzke J, Engels V, Hasenbein S, Gatzens C, Bott M (2008) Coordinated regulation of gluconate catabolism and glucose uptake in *Corynebacterium glutamicum* by two functionally equivalent transcriptional regulators, GntR1 and GntR2. *Mol Microbiol* 67(2):305–322. <https://doi.org/10.1111/j.1365-2958.2007.06020.x>
7. Görke B, Stülke J (2008) Carbon catabolite repression in bacteria: many ways to make the most out of nutrients. *Nat Rev Microbiol* 6(8):613. <https://doi.org/10.1038/nrmicro1932>
8. Rojo F (2010) Carbon catabolite repression in *Pseudomonas*: optimizing metabolic versatility and interactions with the environment. *FEMS Microbiol Rev* 34(5):658–684. <https://doi.org/10.1111/j.1574-6976.2010.00218.x>
9. Collier DN, Hager PW, Phibbs PV (1996) Catabolite repression control in the Pseudomonads. *Res Microbiol* 147(6):551–561. [https://doi.org/10.1016/0923-2508\(96\)84011-3](https://doi.org/10.1016/0923-2508(96)84011-3)
10. Durica-Mitic S, Gopel Y, Gorke B (2018) Carbohydrate utilization in bacteria: making the most out of sugars with the help of small regulatory RNAs. *Microbiol Spectr* 6(2):1–19. <https://doi.org/10.1128/microbiolspec.rwr-0013-2017>
11. Galinier A, Deutscher J (2017) Sophisticated regulation of transcriptional factors by the bacterial phosphoenolpyruvate: sugar phosphotransferase system. *J Mol Biol* 429(6):773–789. <https://doi.org/10.1016/j.jmb.2017.02.006>
12. Vinuselvi P, Kim MK, Lee SK, Ghim CM (2012) Rewiring carbon catabolite repression for microbial cell factory. *BMB Rep* 45(2):59–70. <https://doi.org/10.5483/bmbrep.2012.45.2.59>
13. Monod J (1949) The growth of bacterial cultures. *Annu Rev Microbiol* 3(1):371–394. <https://doi.org/10.1146/annurev.mi.03.100149.002103>
14. Chu D (2017) Limited by sensing—A minimal stochastic model of the lag-phase during diauxic growth. *J Theor Biol* 414:137–146. <https://doi.org/10.1093/bjc/azq038>
15. Chu DF (2015) In silico evolution of diauxic growth. *BMC Evol Biol* 15(1):211. <https://doi.org/10.1186/s12862-015-0492-0>
16. Hermsen R, Okano H, You C, Werner N, Hwa T (2015) A growth-rate composition formula for the growth of *E coli* on co-utilized carbon substrates. *Mol Syst Biol* 11(4):801. <https://doi.org/10.15252/msb.20145537>
17. Solopova A, van Gestel J, Weissing FJ, Bachmann H, Teusink B, Kok J, Kuipers OP (2014) Bet-hedging during bacterial diauxic shift. *Proc Natl Acad Sci*. <https://doi.org/10.1073/pnas.1320063111>
18. Kotte O, Volkmer B, Radzikowski JL, Heinemann M (2014) Phenotypic bistability in *Escherichia coli*'s central carbon metabolism. *Mol Syst Biol* 10:736. <https://doi.org/10.15252/msb.20135022>
19. Succurro A, Segre D, Ebenhoeh O (2019) Emergent subpopulation behavior uncovered with a community dynamic metabolic model of *Escherichia coli* diauxic growth. *mSystems*. <https://doi.org/10.1128/msystems.00230-18>
20. Zimmermann T, Sorg T, Siehler SY, Gerischer U (2009) Role of *Acinetobacter baylyi* Crc in catabolite repression of enzymes for aromatic compound catabolism. *J Bacteriol* 191(8):2834–2842. <https://doi.org/10.1128/JB.00817-08>
21. Cornforth DM, Foster KR (2013) Competition sensing: the social side of bacterial stress responses. *Nat Rev Microbiol* 11:285. <https://doi.org/10.1038/nrmicro2977>
22. Liu W, Roder HL, Madsen JS, Bjarnsholt T, Sorensen SJ, Burmolle M (2016) Interspecific bacterial interactions are reflected in multispecies biofilm spatial organization. *Front Microbiol* 7:1366. <https://doi.org/10.3389/fmicb.2016.01366>
23. Ren D, Madsen JS, Sorensen SJ, Burmolle M (2015) High prevalence of biofilm synergy among bacterial soil isolates in cocultures indicates bacterial interspecific cooperation. *ISME J* 9(1):81–89. <https://doi.org/10.1038/ismej.2014.96>
24. Russel J, Roder HL, Madsen JS, Burmolle M, Sorensen SJ (2017) Antagonism correlates with metabolic similarity in diverse bacteria. *Proc Natl Acad Sci USA* 114(40):10684–10688. <https://doi.org/10.1073/pnas.1706016114>
25. Bernstein HC, Carlson RP (2012) Microbial consortia engineering for cellular factories: in vitro to in silico systems. *Comput Struct Biotechnol J* 3(4):e201210017. <https://doi.org/10.5936/CSBJ.201210017>
26. Scitable by Nature EDUCATION (2005) Nature Publishing Group. www.nature.com/scitable/definition/catabolite-repression-1113/. Accessed 3 Sep 2018
27. Zhang YH, Lynd LR (2005) Cellulose utilization by *Clostridium thermocellum*: bioenergetics and hydrolysis product assimilation. *Proc Natl Acad Sci USA* 102(20):7321–7325. <https://doi.org/10.1073/pnas.0408734102>
28. Parche S, Jacobs D, Arigoni F, Titgemeyer F, Jankovic I (2006) Lactose-over-glucose preference in *Bifidobacterium longum* NCC2705: glcP, encoding a glucose transporter, is subject to lactose repression. *Society* 188(4):1260–1265. <https://doi.org/10.1128/JB.188.4.1260>
29. Ng TK, Zeikus JG (1982) Differential metabolism of cellobiose and glucose by *Clostridium thermocellum* and *Clostridium thermohydrosulfuricum*. *J Bacteriol* 150(3):1391–1399
30. Frimmersdorf E, Horatzek S, Pelnikovich A, Wiehlmann L, Schomburg D (2010) How *Pseudomonas aeruginosa* adapts to various environments: a metabolomic approach. *Environ Microbiol* 12(6):1734–1747. <https://doi.org/10.1111/j.1462-2920.2010.02253.x>
31. Carlson RP (2007) Metabolic systems cost-benefit analysis for interpreting network structure and regulation. *Bioinformatics* 23(10):1258–1264. <https://doi.org/10.1093/bioinformatics/btm082>
32. Carlson RP (2009) Decomposition of complex microbial behaviors into resource-based stress responses. *Bioinformatics* 25(1):90–97
33. Hamilton WA, Dawes EA (1959) A diauxic effect with *Pseudomonas aeruginosa*. *Biochem J* 71:25P–26P
34. Harwood JP, Gazdar C, Prasad C, Peterkofsky A (1976) A involvement of the glucose enzymes II of the sugar phosphotransferase system in the regulation of adenylate cyclase by glucose in *Escherichia coli*. *J Biol Chem* 251(8):2462–2468
35. Feucht BU, Saier MH (1980) Fine control of adenylate cyclase by the phosphoenolpyruvate:sugar phosphotransferase systems in *Escherichia coli* and *Salmonella typhimurium*. *J Bacteriol* 141(2):603–610
36. Titgemeyer F, Hillen W (2002) Global control of sugar metabolism: a Gram-positive solution. *Antonie Van Leeuwenhoek* 82(1):59–71. <https://doi.org/10.1023/A:1020628909429>
37. Warner JB, Lolkema JS (2003) CcpA-dependent carbon catabolite repression in bacteria. *Microbiol Mol Biol Rev* 67(4):475–490. <https://doi.org/10.1128/MMBR.67.4.475-490.2003>
38. Henkin TM, Grundy FJ, Nicholson WL, Chambliss GH (1991) Catabolite repression of α amylase gene expression in *Bacillus subtilis* involves a trans-acting gene product homologous to the *Escherichia coli* lacI and galR repressors. *Mol Microbiol* 5(3):575–584. <https://doi.org/10.1111/j.1365-2958.1991.tb00728.x>
39. Djordjevic GM, Tchiew JH, Saier MH (2001) Genes involved in control of galactose uptake in *Lactobacillus brevis* and reconstitution of the regulatory system in *Bacillus subtilis*. *J Bacteriol* 183(10):3224–3236. <https://doi.org/10.1128/JB.183.10.3224-3236.2001>

40. Zomer AL, Buist G, Larsen R, Kok J, Kuipers OP (2007) Time-resolved determination of the CcpA regulon of *Lactococcus lactis* subsp. *cremoris* MG1363. *J Bacteriol* 189(4):1366–1381. <https://doi.org/10.1128/jb.01013-06>
41. Abdou L, Boileau C, Philip Pd, Pagès S, Fiérobe H-P, Tardif C (2008) Transcriptional regulation of the clostridium cellulolyticum cip-cel Operon: a complex mechanism involving a catabolite-responsive element. *J Bacteriol* 190(5):1499–1506. <https://doi.org/10.1128/JB.01160-07>
42. Arndt A, Eikmanns BJ (2007) The alcohol dehydrogenase gene adhA in corynebacterium glutamicum is subject to carbon catabolite repression. *J Bacteriol* 189(20):7408–7416. <https://doi.org/10.1128/JB.00791-07>
43. Gerstmeir R, Cramer A, Dangel P, Schaffer S, Eikmanns BJ (2004) RamB, a novel transcriptional regulator of genes involved in acetate metabolism of *Corynebacterium glutamicum*. *J Bacteriol* 186(9):2798–2809
44. Müller C, Petruschka L, Cuypers H, Burchhardt G, Herrmann H (1996) Carbon catabolite repression of phenol degradation in *Pseudomonas putida* is mediated by the inhibition of the activator protein PhlR. *J Bacteriol* 178(7):2030–2036. <https://doi.org/10.1128/jb.178.7.2030-2036.1996>
45. Kwakman JH, Postma PW (1994) Glucose kinase has a regulatory role in carbon catabolite repression in *Streptomyces coelicolor*. *J Bacteriol* 176(9):2694–2698. <https://doi.org/10.1128/jb.176.9.2694-2698.1994>
46. Commichau FM, Stülke J (2008) Trigger enzymes: bifunctional proteins active in metabolism and in controlling gene expression. *Mol Microbiol* 67(4):692–702. <https://doi.org/10.1111/j.1365-2958.2007.06071.x>
47. Chu D, Barnes DJ (2016) The lag-phase during diauxic growth is a trade-off between fast adaptation and high growth rate. *Sci Rep* 6:25191. <https://doi.org/10.1038/srep25191>
48. Wezel GPV, König M, Mahr K, Nothaft H, Thomae AW, Bibb M, Titgemeyer F (2007) A new piece of an old jigsaw: glucose kinase is activated posttranslationally in a glucose transport-dependent manner in streptomyces coelicolor A3(2). *MMB* 12(1–2):67–74. <https://doi.org/10.1159/000096461>
49. Busby S, Ebright RH (1999) Transcription activation by catabolite activator protein (CAP). *J Mol Biol* 293(2):199–213. <https://doi.org/10.1006/jmbi.1999.3161>
50. Malan TP, Kolb A, Buc H, McClure WR (1984) Mechanism of CRP-cAMP activation of lac operon transcription initiation activation of the P1 promoter. *J Mol Biol* 180(4):881–909. [https://doi.org/10.1016/0022-2836\(84\)90262-6](https://doi.org/10.1016/0022-2836(84)90262-6)
51. Deutscher J (2008) The mechanisms of carbon catabolite repression in bacteria. *Curr Opin Microbiol* 11(2):87–93. <https://doi.org/10.1016/j.mib.2008.02.007>
52. Harwood JP, Gazdar C, Prasad C, Peterkofsky A, Curtis SJ, Epstein W (1976) Involvement of the glucose enzymes II of the sugar phosphotransferase system in the regulation of adenylate cyclase by glucose in *Escherichia coli*. *J Biol Chem* 251(8):2462–2468
53. Park Y-H, Lee BR, Seok Y-J, Peterkofsky A (2006) In vitro reconstitution of catabolite repression in *Escherichia coli*. *J Biol Chem* 281(10):6448–6454. <https://doi.org/10.1074/jbc.M512672200>
54. Tagami H, Aiba H (1998) A common role of CRP in transcription activation: CRP acts transiently to stimulate events leading to open complex formation at a diverse set of promoters. *EMBO J* 17(6):1759–1767. <https://doi.org/10.1093/emboj/17.6.1759>
55. Cases I, Velázquez F, de Lorenzo V (2007) The ancestral role of the phosphoenolpyruvate–carbohydrate phosphotransferase system (PTS) as exposed by comparative genomics. *Res Microbiol* 158(8):666–670. <https://doi.org/10.1016/j.resmic.2007.08.002>
56. Deutscher J, Francke C, Postma PW (2006) How phosphotransferase system-related protein phosphorylation regulates carbohydrate metabolism in bacteria. *Microbiol Mol Biol Rev* 70(4):939–1031. <https://doi.org/10.1128/MMBR.00024-06>
57. Postma PW, Lengeler JW, Jacobson GR (1993) Phosphoenolpyruvate:carbohydrate phosphotransferase systems of bacteria. *Microbiol Mol Biol Rev* 57(3):543–594
58. Hogema BM, Arents JC, Bader R, Postma PW (1999) Autoregulation of lactose uptake through the LacY permease by enzyme IIAGlc of the PTS in *Escherichia coli* K-12. *Mol Microbiol* 31(6):1825–1833. <https://doi.org/10.1046/j.1365-2958.1999.01319.x>
59. Inada T, Kimata K, Aiba H (1996) Mechanism responsible for glucose–lactose diauxie in *Escherichia coli*: challenge to the cAMP model. *Genes Cells* 1(3):293–301. <https://doi.org/10.1046/j.1365-2443.1996.24025.x>
60. Kimata K, Takahashi H, Inada T, Postma P, Aiba H (1997) cAMP receptor protein–cAMP plays a crucial role in glucose–lactose diauxie by activating the major glucose transporter gene in *Escherichia coli*. *Proc Natl Acad Sci* 94(24):12914–12919. <https://doi.org/10.1073/pnas.94.24.12914>
61. Beckwith JR (1967) Regulation of the Lac operon. *Science* 156(3775):597–604
62. Storz G, Vogel J, Wassarman KM (2011) Regulation by small RNAs in bacteria: expanding frontiers. *Mol Cell* 43(6):880–891. <https://doi.org/10.1016/j.molcel.2011.08.022>
63. Beisel CL, Updegrove TB, Janson BJ, Storz G (2012) Multiple factors dictate target selection by Hfq-binding small RNAs. *EMBO J* 31(8):1961–1974. <https://doi.org/10.1038/emboj.2012.52>
64. Beisel CL, Storz G (2011) Discriminating tastes: physiological contributions of the Hfq-binding small RNA Spot 42 to catabolite repression. *RNA Biol* 8(5):766–770. <https://doi.org/10.4161/rna.8.5.16024>
65. Beisel CL, Storz G (2011) The base-pairing RNA spot 42 participates in a multioutput feedforward loop to help enact catabolite repression in *Escherichia coli*. *Mol Cell* 41(3):286–297. <https://doi.org/10.1016/j.molcel.2010.12.027>
66. You C, Okano H, Hui S, Zhang Z, Kim M, Gunderson CW, Wang YP, Lenz P, Yan D, Hwa T (2013) Coordination of bacterial proteome with metabolism by cyclic AMP signalling. *Nature* 500(7462):301–306. <https://doi.org/10.1038/nature12446>
67. Bettenbrock K, Fischer S, Kremling A, Jahreis K, Sauter T, Gilles ED (2006) A quantitative approach to catabolite repression in *Escherichia coli*. *J Biol Chem* 281(5):2578–2584. <https://doi.org/10.1074/jbc.M508090200>
68. Bren A, Park JO, Towbin BD, Dekel E, Rabinowitz JD, Alon U (2016) Glucose becomes one of the worst carbon sources for *E. coli* on poor nitrogen sources due to suboptimal levels of cAMP. *Sci Rep* 6:24834. <https://doi.org/10.1038/srep24834>
69. Deutscher J, Küster E, Bergstedt U, Charrier V, Hillen W (1995) Protein kinase-dependent HPr/CcpA interaction links glycolytic activity to carbon catabolite repression in Gram-positive bacteria. *Mol Microbiol* 15(6):1049–1053. <https://doi.org/10.1111/j.1365-2958.1995.tb02280.x>
70. Jones BE, Dossonnet V, Küster E, Hillen W, Deutscher J, Klevit RE (1997) Binding of the catabolite repressor protein CcpA to its DNA target is regulated by phosphorylation of its corepressor HPr. *J Biol Chem* 272(42):26530–26535. <https://doi.org/10.1074/jbc.272.42.26530>
71. Galinier A, Kravanja M, Engelmann R, Hengstenberg W, Kilhofer M-C, Deutscher J, Haiech J (1998) New protein kinase and protein phosphatase families mediate signal transduction in bacterial catabolite repression. *Proc Natl Acad Sci* 95(4):1823–1828. <https://doi.org/10.1073/pnas.95.4.1823>

72. Jault J-M, Fieulaine S, Nessler S, Gonzalo P, Pietro AD, Deutscher J, Galinier A (2000) The HPr kinase from *Bacillus subtilis* is a homo-oligomeric enzyme which exhibits strong positive cooperativity for nucleotide and fructose 1,6-bisphosphate binding. *J Biol Chem* 275(3):1773–1780. <https://doi.org/10.1074/jbc.275.3.1773>
73. Reizer J, Hoischen C, Titgemeyer F, Rivolta C, Rabus R, Stülke J, Karamata D, Hillen W (1998) A novel protein kinase that controls carbon catabolite repression in bacteria. *Mol Microbiol* 27(6):1157–1169. <https://doi.org/10.1046/j.1365-2958.1998.00747.x>
74. Schumacher MA, Allen GS, Diel M, Seidel G, Hillen W, Brennan RG (2004) Structural basis for allosteric control of the transcription regulator CcpA by the phosphoprotein HPr-Ser46-P. *Cell* 118(6):731–741. <https://doi.org/10.1016/j.cell.2004.08.027>
75. Seidel G, Diel M, Fuchsbauer N, Hillen W (2005) Quantitative interdependence of coeffectors, CcpA and cre in carbon catabolite regulation of *Bacillus subtilis*. *FEBS J* 272(10):2566–2577. <https://doi.org/10.1111/j.1742-4658.2005.04682.x>
76. Moreno MS, Schneider BL, Maile RR, Weyler W, Saier MH (2004) Catabolite repression mediated by the CcpA protein in *Bacillus subtilis*: novel modes of regulation revealed by whole-genome analyses. *Mol Microbiol* 39(5):1366–1381. <https://doi.org/10.1111/j.1365-2958.2001.02328.x>
77. Yoshida K, Kobayashi K, Miwa Y, Kang CM, Matsunaga M, Yamaguchi H, Tojo S, Yamamoto M, Nishi R, Ogasawara N, Nakayama T, Fujita Y (2001) Combined transcriptome and proteome analysis as a powerful approach to study genes under glucose repression in *Bacillus subtilis*. *Nucleic Acids Res* 29(3):683–692. <https://doi.org/10.1093/nar/29.3.683>
78. Kim SU, Kim DC, Dhurjati P (1988) Mathematical modeling for mixed culture growth of two bacterial populations with opposite substrate preferences. *Biotechnol Bioeng* 31(2):144–159. <https://doi.org/10.1002/bit.260310208>
79. Ramkrishna D, Kompala DS, Tsao GT (1987) Are microbes optimal strategists? *Biotechnol Progr* 3(3):121–126. <https://doi.org/10.1002/btpr.5420030302>
80. Kim SU, Dhurjati P (1987) Analysis of two interacting bacterial populations with opposite substrate preferences. *Biotechnol Bioeng* 29(8):1015–1023. <https://doi.org/10.1002/bit.260290813>
81. Wolff JA, MacGregor CH, Eisenberg RC, Phibbs PV (1991) Isolation and characterization of catabolite repression control mutants of *Pseudomonas aeruginosa* PAO. *J Bacteriol* 173(15):4700–4706
82. Stewart PS (2003) Diffusion in Biofilms. *J Bacteriol* 185(5):1485–1491. <https://doi.org/10.1128/JB.185.5.1485-1491.2003>
83. Sonnleitner E, Blasi U (2014) Regulation of Hfq by the RNA CrcZ in *Pseudomonas aeruginosa* carbon catabolite repression. *PLoS Genet* 10(6):e1004440. <https://doi.org/10.1371/journal.pgen.1004440>
84. Sonnleitner E, Abdou L, Haas D (2009) Small RNA as global regulator of carbon catabolite repression in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci* 106(51):21866–21871. <https://doi.org/10.1073/pnas.0910308106>
85. Sonnleitner E, Wulf A, Campagne S, Pei XY, Wolfinger MT, Forlani G, Prindl K, Abdou L, Resch A, Allain FH, Luisi BF, Urlaub H, Blasi U (2018) Interplay between the catabolite repression control protein Crc, Hfq and RNA in Hfq-dependent translational regulation in *Pseudomonas aeruginosa*. *Nucleic Acids Res* 46(3):1470–1485. <https://doi.org/10.1093/nar/gkx1245>
86. Moreno R, Hernandez-Arranz S, La Rosa R, Yuste L, Madhushani A, Shingler V, Rojo F (2015) The Crc and Hfq proteins of *Pseudomonas putida* cooperate in catabolite repression and formation of ribonucleic acid complexes with specific target motifs. *Environ Microbiol* 17(1):105–118. <https://doi.org/10.1111/1462-2920.12499>
87. Kambara TK, Ramsey KM, Dove SL (2018) Pervasive targeting of nascent transcripts by Hfq. *Cell Rep* 23(5):1543–1552. <https://doi.org/10.1016/j.celrep.2018.03.134>
88. Grenga L, Chandra G, Saalbach G, Galmozzi CV, Kramer G, Malone JG (2017) Analyzing the complex regulatory landscape of Hfq—an integrative, multi-omics approach. *Front Microbiol* 8:1784. <https://doi.org/10.3389/fmicb.2017.01784>
89. Moreno R, Fonseca P, Rojo F (2012) Two small RNAs, CrcY and CrcZ, act in concert to sequester the Crc global regulator in *Pseudomonas putida*, modulating catabolite repression. *Mol Microbiol* 83(1):24–40. <https://doi.org/10.1111/j.1365-2958.2011.07912.x>
90. Liu Y, Gokhale CS, Rainey PB, Zhang XX (2017) Unravelling the complexity and redundancy of carbon catabolic repression in *Pseudomonas fluorescens* SBW25. *Mol Microbiol* 105(4):589–605. <https://doi.org/10.1111/mmi.13720>
91. Filiatrault MJ, Stodghill PV, Wilson J, Butcher BG, Chen H, Myers CR, Cartinhour SW (2013) CrcZ and CrcX regulate carbon source utilization in *Pseudomonas syringae* pathovar tomato strain DC3000. *RNA Biol* 10(2):245–255. <https://doi.org/10.4161/rna.23019>
92. Garcia-Maurino SM, Perez-Martinez I, Amador CI, Canosa I, Santero E (2013) Transcriptional activation of the CrcZ and CrcY regulatory RNAs by the CbrB response regulator in *Pseudomonas putida*. *Mol Microbiol* 89(1):189–205. <https://doi.org/10.1111/mmi.12270>
93. Kremling A, Geiselmann J, Ropers D, de Jong H (2015) Understanding carbon catabolite repression in *Escherichia coli* using quantitative models. *Trends Microbiol* 23(2):99–109. <https://doi.org/10.1016/j.tim.2014.11.002>
94. Kremling A, Geiselmann J, Ropers D, de Jong H (2018) An ensemble of mathematical models showing diauxic growth behaviour. *BMC Syst Biol* 12(1):1–16. <https://doi.org/10.1186/s12918-018-0604-8>
95. Kompala DS, Ramkrishna D, Jansen NB, Tsao GT (1986) Investigation of bacterial growth on mixed substrates: experimental evaluation of cybernetic models. *Biotechnol Bioeng* 28(7):1044–1055. <https://doi.org/10.1002/bit.260280715>
96. DeVilbiss F, Mandli A, Ramkrishna D (2018) Consistency of cybernetic variables with gene expression profiles: a more rigorous test. *Biotechnol Prog* 34(4):858–867. <https://doi.org/10.1002/btpr.2654>
97. Schuster R, Schuster S (1993) Refined algorithm and computer program for calculating all non-negative fluxes admissible in steady states of biochemical reaction systems with or without some flux rates fixed. *Bioinformatics* 9(1):79–85. <https://doi.org/10.1093/bioinformatics/9.1.79>
98. Schuster S, Dandekar T, Fell DA (1999) Detection of elementary flux modes in biochemical networks: a promising tool for pathway analysis and metabolic engineering. *Trends Biotechnol* 17(2):53–60. [https://doi.org/10.1016/S0167-7799\(98\)01290-6](https://doi.org/10.1016/S0167-7799(98)01290-6)
99. Varma A, Boesch BW, Palsson BO (1993) Stoichiometric interpretation of *Escherichia coli* glucose catabolism under various oxygenation rates. *Appl Environ Microbiol* 59(8):2465–2473
100. Varma A, Palsson BO (1994) Stoichiometric flux balance models quantitatively predict growth and metabolic by-product secretion in wild-type *Escherichia-Coli* W3110. *Appl Environ Microb* 60(10):3724–3731
101. Beck AE, Hunt KA, Carlson RP (2018) Measuring cellular biomass composition for computational biology applications. *Processes* 6(5):38. <https://doi.org/10.3390/pr6050038>
102. Schuster S, Pfeiffer T, Fell DA (2008) Is maximization of molar yield in metabolic networks favoured by evolution? *J Theor Biol* 252(3):497–504. <https://doi.org/10.1016/j.jtbi.2007.12.008>
103. Beg QK, Vazquez A, Ernst J, de Menezes MA, Bar-Joseph Z, Barabasi AL, Oltvai ZN (2007) Intracellular crowding defines

- the mode and sequence of substrate uptake by *Escherichia coli* and constrains its metabolic activity. *Proc Natl Acad Sci USA* 104(31):12663–12668. <https://doi.org/10.1073/pnas.0609845104>
104. Vazquez A, Oltvai ZN (2016) Macromolecular crowding explains overflow metabolism in cells. *Sci Rep* 6:31007. <https://doi.org/10.1038/srep31007>
 105. Carlson R, Sreenc F (2004) Fundamental *Escherichia coli* biochemical pathways for biomass and energy production: identification of reactions. *Biotechnol Bioeng* 85(1):1–19. <https://doi.org/10.1002/bit.10812>
 106. Carlson R, Sreenc F (2004) Fundamental *Escherichia coli* biochemical pathways for biomass and energy production: creation of overall flux states. *Biotechnol Bioeng* 86(2):149–162. <https://doi.org/10.1002/bit.20044>
 107. Carlson RP, Beck AE, Phalak P, Fields MW, Gedeon T, Hanley L, Harcombe WR, Henson MA, Heys JJ (2018) Competitive resource allocation to metabolic pathways contributes to overflow metabolisms and emergent properties in cross-feeding microbial consortia. *Biochem Soc Trans* 46(2):269–284. <https://doi.org/10.1042/BST20170242>
 108. Zhuang K, Vemuri GN, Mahadevan R (2011) Economics of membrane occupancy and respiration-fermentation. *Mol Syst Biol* 7:500. <https://doi.org/10.1038/msb.2011.34>
 109. Szenk M, Dill KA, de Graff AMR (2017) Why do fast-growing bacteria enter overflow metabolism? Testing the membrane real estate hypothesis. *Cell Syst* 5(2):95–104. <https://doi.org/10.1016/j.cels.2017.06.005>
 110. Basan M, Hui S, Okano H, Zhang Z, Shen Y, Williamson JR, Hwa T (2015) Overflow metabolism in *Escherichia coli* results from efficient proteome allocation. *Nature* 528(7580):99–104. <https://doi.org/10.1038/nature15765>
 111. Goelzer A (1810) Fromion V (2011) Bacterial growth rate reflects a bottleneck in resource allocation. *Biochim Biophys Acta* 10:978–988. <https://doi.org/10.1016/j.bbagen.2011.05.014>
 112. Goelzer A, Fromion V (2017) Resource allocation in living organisms. *Biochem Soc Trans* 45(4):945–952. <https://doi.org/10.1042/BST20160436>
 113. Yang L, Ma D, Ebrahim A, Lloyd CJ, Saunders MA, Palsson BO (2016) solveME: fast and reliable solution of nonlinear ME models. *BMC Bioinformatics* 17(1):391. <https://doi.org/10.1186/s12859-016-1240-1>
 114. Mahadevan R, Edwards JS, Doyle FJ (2002) Dynamic flux balance analysis of diauxic growth in *Escherichia coli*. *Biophys J* 83(3):1331–1340
 115. Rentz JA, Alvarez PJ, Schnoor JL (2004) Repression of *Pseudomonas putida* phenanthrene-degrading activity by plant root extracts and exudates. *Environ Microbiol* 6(6):574–583. <https://doi.org/10.1111/j.1462-2920.2004.00589.x>
 116. Tiwari N, Campbell J (1969) Enzymatic control of the metabolic activity of *Pseudomonas aeruginosa* grown in glucose or succinate media. *Biochim Biophys Acta (BBA) Gen Subj* 192(3):395–401. [https://doi.org/10.1016/0304-4165\(69\)90388-2](https://doi.org/10.1016/0304-4165(69)90388-2)
 117. Trautwein K, Grundmann O, Wohlbrand L, Eberlein C, Boll M, Rabus R (2012) Benzoate mediates repression of C(4)-dicarboxylate utilization in “*Aromatoleum aromaticum*” EbN1. *J Bacteriol* 194(2):518–528. <https://doi.org/10.1128/JB.05072-11>
 118. Berger A, Dohnt K, Tielen P, Jahn D, Becker J, Wittmann C (2014) Robustness and plasticity of metabolic pathway flux among uropathogenic isolates of *Pseudomonas aeruginosa*. *PLoS One* 9(4):e88368. <https://doi.org/10.1371/journal.pone.0088368>
 119. Hintermayer SB, Weuster-Botz D (2017) Experimental validation of in silico estimated biomass yields of *Pseudomonas putida* KT2440. *Biotechnol J* 12(6):1600720. <https://doi.org/10.1002/biot.201600720>
 120. La Rosa R, Behrends V, Williams HD, Bundy JG, Rojo F (2016) Influence of the Crc regulator on the hierarchical use of carbon sources from a complete medium in *Pseudomonas*. *Environ Microbiol* 18(3):807–818. <https://doi.org/10.1111/1462-2920.13126>
 121. Nikel PI, Chavarria M, Fuhrer T, Sauer U, de Lorenzo V (2015) *Pseudomonas putida* KT2440 strain metabolizes glucose through a cycle formed by enzymes of the entner-doudoroff, embden-meyerhof-parnas, and pentose phosphate pathways. *J Biol Chem* 290(43):25920–25932. <https://doi.org/10.1074/jbc.M115.687749>
 122. Flamholz A, Noor E, Bar-Even A, Liebermeister W, Milo R (2013) Glycolytic strategy as a tradeoff between energy yield and protein cost. *Proc Natl Acad Sci USA* 110(24):10039–10044. <https://doi.org/10.1073/pnas.1215283110>
 123. Oberhardt MA, Goldberg JB, Hogardt M, Papin JA (2010) Metabolic network analysis of *Pseudomonas aeruginosa* during chronic cystic fibrosis lung infection. *J Bacteriol* 192(20):5534–5548. <https://doi.org/10.1128/JB.00900-10>
 124. Xu Z, Fang X, Wood TK, Huang ZJ (2013) A systems-level approach for investigating *Pseudomonas aeruginosa* biofilm formation. *PLoS One* 8(2):e57050. <https://doi.org/10.1371/journal.pone.0057050>
 125. Nogales J, Palsson BO, Thiele I (2008) A genome-scale metabolic reconstruction of *Pseudomonas putida* KT2440: iJN746 as a cell factory. *BMC Syst Biol* 2:79. <https://doi.org/10.1186/1752-0509-2-79>
 126. Borgos SE, Bordel S, Sletta H, Ertesvag H, Jakobsen O, Bruheim P, Ellingsen TE, Nielsen J, Valla S (2013) Mapping global effects of the anti-sigma factor MucA in *Pseudomonas fluorescens* SBW25 through genome-scale metabolic modeling. *BMC Syst Biol* 7:19. <https://doi.org/10.1186/1752-0509-7-19>
 127. Stolyar S, Van Dien S, Hillesland KL, Pinel N, Lie TJ, Leigh JA, Stahl DA (2007) Metabolic modeling of a mutualistic microbial community. *Mol Syst Biol* 3:92. <https://doi.org/10.1038/msb4100131>
 128. Goyal A, Dubinkina V, Maslov S (2018) Multiple stable states in microbial communities explained by the stable marriage problem. *ISME J* 12(12):2823–2834. <https://doi.org/10.1038/s41396-018-0222-x>
 129. Roell GW, Zha J, Carr RR, Koffas MA, Fong SS, Tang YJ (2019) Engineering microbial consortia by division of labor. *Microb Cell Fact* 18(1):35. <https://doi.org/10.1186/s12934-019-1083-3>
 130. Shuler ML, Kargi F, DeLis M (2017) Bioprocess engineering: basic concepts, 3rd edn. Prentice Hall, Boston
 131. Kreft JU (2004) Biofilms promote altruism. *Microbiology* 150(Pt 8):2751–2760. <https://doi.org/10.1099/mic.0.26829-0>
 132. Boyle KE, Monaco HT, Deforet M, Yan J, Wang Z, Rhee K, Xavier JB (2017) Metabolism and the evolution of social behavior. *Mol Biol Evol* 34(9):2367–2379. <https://doi.org/10.1093/molbev/msx174>
 133. Ng WL, Bassler BL (2009) Bacterial quorum-sensing network architectures. *Annu Rev Genet* 43:197–222. <https://doi.org/10.1146/annurev-genet-102108-134304>
 134. Sonnleitner E, Schuster M, Sorger-Domenigg T, Greenberg EP, Blasi U (2006) Hfq-dependent alterations of the transcriptome profile and effects on quorum sensing in *Pseudomonas aeruginosa*. *Mol Microbiol* 59(5):1542–1558. <https://doi.org/10.1111/j.1365-2958.2006.05032.x>
 135. Zhang L, Gao Q, Chen W, Qin H, Hengzhuang W, Chen Y, Yang L, Zhang G (2013) Regulation of pqs quorum sensing via catabolite repression control in *Pseudomonas aeruginosa*. *Microbiology* 159(Pt 9):1931–1936. <https://doi.org/10.1099/mic.0.066266-0>
 136. Yang N, Lan L (2016) *Pseudomonas aeruginosa* Lon and ClpXP proteases: roles in linking carbon catabolite repression system

- with quorum-sensing system. *Curr Genet* 62(1):1–6. <https://doi.org/10.1007/s00294-015-0499-5>
137. Parsek MR, Greenberg EP (2005) Sociomicrobiology: the connections between quorum sensing and biofilms. *Trends Microbiol* 13(1):27–33. <https://doi.org/10.1016/j.tim.2004.11.007>
 138. Coenye T (2010) Social interactions in the *Burkholderia cepacia* complex: biofilms and quorum sensing. *Future Microbiol* 5(7):1087–1099. <https://doi.org/10.2217/fmb.10.68>
 139. Hall-Stoodley L, Costerton JW, Stoodley P (2004) Bacterial biofilms: from the natural environment to infectious diseases. *Nat Rev Microbiol* 2(2):95–108. <https://doi.org/10.1038/nrmicro821>
 140. Flemming HC, Wingender J, Szewzyk U, Steinberg P, Rice SA, Kjelleberg S (2016) Biofilms: an emergent form of bacterial life. *Nat Rev Microbiol* 14(9):563–575. <https://doi.org/10.1038/nrmicro.2016.94>
 141. Xavier JB, Foster KR (2007) Cooperation and conflict in microbial biofilms. *Proc Natl Acad Sci USA* 104(3):876–881. <https://doi.org/10.1073/pnas.0607651104>
 142. Nadell CD, Drescher K, Foster KR (2016) Spatial structure, cooperation and competition in biofilms. *Nat Rev Microbiol* 14(9):589–600. <https://doi.org/10.1038/nrmicro.2016.84>
 143. Stewart PS (1998) A review of experimental measurements of effective diffusive permeabilities and effective diffusion coefficients in biofilms. *Biotechnol Bioeng* 59(3):261–272. [https://doi.org/10.1002/\(SICI\)1097-0290\(19980805\)59:3%3c261::AID-BIT1%3e3.0.CO;2-9](https://doi.org/10.1002/(SICI)1097-0290(19980805)59:3%3c261::AID-BIT1%3e3.0.CO;2-9)
 144. Kreft JU, Bonhoeffer S (2005) The evolution of groups of cooperating bacteria and the growth rate versus yield trade-off. *Microbiology* 151(Pt 3):637–641. <https://doi.org/10.1099/mic.0.27415-0>
 145. Liu J, Prindle A, Humphries J, Gabalda-Sagarra M, Asally M, Lee DY, Ly S, Garcia-Ojalvo J, Suel GM (2015) Metabolic co-dependence gives rise to collective oscillations within biofilms. *Nature* 523(7562):550–554. <https://doi.org/10.1038/nature14660>
 146. Liu J, Martinez-Corral R, Prindle A, Lee DD, Larkin J, Gabalda-Sagarra M, Garcia-Ojalvo J, Suel GM (2017) Coupling between distant biofilms and emergence of nutrient time-sharing. *Science* 356(6338):638–642. <https://doi.org/10.1126/science.aah4204>
 147. Johnson DI (2018) *Pseudomonas* spp. In: Bacterial pathogens and their virulence factors, Springer International Publishing, Cham, pp 325–338. https://doi.org/10.1007/978-3-319-67651-7_25
 148. Pusic P, Sonnleitner E, Krennmayr B, Heitzinger DA, Wolfinger MT, Resch A, Blasi U (2018) Harnessing metabolic regulation to increase Hfq-dependent antibiotic susceptibility in *Pseudomonas aeruginosa*. *Front Microbiol* 9:2709. <https://doi.org/10.3389/fmicb.2018.02709>
 149. Corona F, Martinez JL, Nikel PI (2019) The global regulator Crc orchestrates the metabolic robustness underlying oxidative stress resistance in *Pseudomonas aeruginosa*. *Environ Microbiol* 21(3):898–912. <https://doi.org/10.1111/1462-2920.14471>
 150. Bernal V, Castano-Cerezo S, Canovas M (2016) Acetate metabolism regulation in *Escherichia coli*: carbon overflow, pathogenicity, and beyond. *Appl Microbiol Biotechnol* 100(21):8985–9001. <https://doi.org/10.1007/s00253-016-7832-x>
 151. Kelsic ED, Zhao J, Vetsigian K, Kishony R (2015) Counteraction of antibiotic production and degradation stabilizes microbial communities. *Nature* 521(7553):516–519. <https://doi.org/10.1038/nature14485>
 152. Hibbing ME, Fuqua C, Parsek MR, Peterson SB (2010) Bacterial competition: surviving and thriving in the microbial jungle. *Nat Rev Microbiol* 8(1):15–25. <https://doi.org/10.1038/nrmicro2259>
 153. Meylan S, Porter CBM, Yang JH, Belenky P, Gutierrez A, Lobritz MA, Park J, Kim SH, Moskowitz SM, Collins JJ (2017) carbon sources tune antibiotic susceptibility in *Pseudomonas aeruginosa* via tricarboxylic acid cycle control. *Cell Chem Biol* 24(2):195–206. <https://doi.org/10.1016/j.chembiol.2016.12.015>
 154. Yeung AT, Bains M, Hancock RE (2011) The sensor kinase CbrA is a global regulator that modulates metabolism, virulence, and antibiotic resistance in *Pseudomonas aeruginosa*. *J Bacteriol* 193(4):918–931. <https://doi.org/10.1128/JB.00911-10>
 155. Hufnagel DA, Evans ML, Greene SE, Pinkner JS, Hultgren SJ, Chapman MR (2016) The catabolite repressor protein-cyclic AMP complex regulates csgD and biofilm formation in uropathogenic *Escherichia coli*. *J Bacteriol* 198(24):3329–3334. <https://doi.org/10.1128/JB.00652-16>
 156. Jackson DW, Simecka JW, Romeo T (2002) Catabolite repression of *Escherichia coli* biofilm formation. *J Bacteriol* 184(12):3406–3410. <https://doi.org/10.1128/jb.184.12.3406-3410.2002>
 157. O'Toole GA, Gibbs KA, Hager PW, Phibbs PV Jr, Kolter R (2000) The global carbon metabolism regulator Crc is a component of a signal transduction pathway required for biofilm development by *Pseudomonas aeruginosa*. *J Bacteriol* 182(2):425–431. <https://doi.org/10.1128/jb.182.2.425-431.2000>
 158. Pusic P, Tata M, Wolfinger MT, Sonnleitner E, Haussler S, Blasi U (2016) Cross-regulation by CrcZ RNA controls anoxic biofilm formation in *Pseudomonas aeruginosa*. *Sci Rep* 6:39621. <https://doi.org/10.1038/srep39621>
 159. Amador CI, Lopez-Sanchez A, Govantes F, Santero E, Canosa I (2016) A *Pseudomonas putida* cbrB transposon insertion mutant displays a biofilm hyperproducing phenotype that is resistant to dispersal. *Environ Microbiol Rep* 8(5):622–629. <https://doi.org/10.1111/1758-2229.12414>
 160. Seidl K, Goerke C, Wolz C, Mack D, Berger-Bachi B, Bischoff M (2008) *Staphylococcus aureus* CcpA affects biofilm formation. *Infect Immun* 76(5):2044–2050. <https://doi.org/10.1128/IAI.00035-08>
 161. Sadykov MR, Hartmann T, Mattes TA, Hiatt M, Jann NJ, Zhu Y, Ledala N, Landmann R, Herrmann M, Rohde H, Bischoff M, Somerville GA (2011) CcpA coordinates central metabolism and biofilm formation in *Staphylococcus epidermidis*. *Microbiology* 157(Pt 12):3458–3468. <https://doi.org/10.1099/mic.0.051243-0>
 162. Ha JH, Hauk P, Cho K, Eo Y, Ma X, Stephens K, Cha S, Jeong M, Suh JY, Sintim HO, Bentley WE, Ryu KS (2018) Evidence of link between quorum sensing and sugar metabolism in *Escherichia coli* revealed via cocrystal structures of LsrK and HPr. *Sci Adv* 4(6):eaar7063. <https://doi.org/10.1126/sciadv.aar7063>
 163. Mitra A, Herren CD, Patel IR, Coleman A, Mukhopadhyay S (2016) Integration of AI-2 based cell-cell signaling with metabolic cues in *Escherichia coli*. *PLoS One* 11(6):e0157532. <https://doi.org/10.1371/journal.pone.0157532>
 164. Wang L, Hashimoto Y, Tsao CY, Valdes JJ, Bentley WE (2005) Cyclic AMP (cAMP) and cAMP receptor protein influence both synthesis and uptake of extracellular autoinducer 2 in *Escherichia coli*. *J Bacteriol* 187(6):2066–2076. <https://doi.org/10.1128/JB.187.6.2066-2076.2005>
 165. Novick RP, Geisinger E (2008) Quorum sensing in staphylococci. *Annu Rev Genet* 42:541–564. <https://doi.org/10.1146/annurev.genet.42.110807.091640>
 166. Hartmann T, Baronian G, Nippe N, Voss M, Schulthess B, Wolz C, Eisenbeis J, Schmidt-Hohagen K, Gaupp R, Sunderkotter C, Beisswenger C, Bals R, Somerville GA, Herrmann M, Molle V, Bischoff M (2014) The catabolite control protein E (CcpE) affects virulence determinant production and pathogenesis of *Staphylococcus aureus*. *J Biol Chem* 289(43):29701–29711. <https://doi.org/10.1074/jbc.M114.584979>
 167. Chakravarthy S, Butcher BG, Liu Y, D'Amico K, Coster M, Filiatrault MJ (2017) Virulence of *Pseudomonas syringae* pv. tomato DC3000 is influenced by the catabolite repression control protein Crc. *Mol Plant Microbe Interact* 30(4):283–294. <https://doi.org/10.1094/mpmi-09-16-0196-r>

168. Tuncil YE, Xiao Y, Porter NT, Reuhs BL, Martens EC, Hamaker BR (2017) Reciprocal prioritization to dietary glycans by gut bacteria in a competitive environment promotes stable coexistence. *MBio* 8(5):e01068–17. <https://doi.org/10.1128/mbio.01068-17>
169. Riviere A, Selak M, Geirnaert A, Van den Abbeele P, De Vuyst L (2018) Complementary mechanisms for degradation of inulin-type fructans and arabinoxylan oligosaccharides among bifidobacterial strains suggest bacterial cooperation. *Appl Environ Microbiol*. <https://doi.org/10.1128/AEM.02893-17>
170. Foster KR, Bell T (2012) Competition, not cooperation, dominates interactions among culturable microbial species. *Curr Biol* 22(19):1845–1850. <https://doi.org/10.1016/j.cub.2012.08.005>
171. Orazi G, O'Toole GA (2017) *Pseudomonas aeruginosa* alters *Staphylococcus aureus* sensitivity to vancomycin in a biofilm model of cystic fibrosis infection. *MBio* 8(4). <https://doi.org/10.1128/mbio.00873-17>
172. Rhoads DD, Wolcott RD, Sun Y, Dowd SE (2012) Comparison of culture and molecular identification of bacteria in chronic wounds. *Int J Mol Sci* 13(3):2535–2550. <https://doi.org/10.3390/ijms13032535>
173. Dalton T, Dowd SE, Wolcott RD, Sun Y, Watters C, Griswold JA, Rumbaugh KP (2011) An in vivo polymicrobial biofilm wound infection model to study interspecies interactions. *PLoS One* 6(11):e27317. <https://doi.org/10.1371/journal.pone.0027317>
174. Gjodsbol K, Christensen JJ, Karlsmark T, Jorgensen B, Klein BM, Krogfelt KA (2006) Multiple bacterial species reside in chronic wounds: a longitudinal study. *Int Wound J* 3(3):225–231. <https://doi.org/10.1111/j.1742-481X.2006.00159.x>
175. Kirketerp-Moller K, Jensen PO, Fazli M, Madsen KG, Pedersen J, Moser C, Tolker-Nielsen T, Hoiby N, Givskov M, Bjarnsholt T (2008) Distribution, organization, and ecology of bacteria in chronic wounds. *J Clin Microbiol* 46(8):2717–2722. <https://doi.org/10.1128/JCM.00501-08>
176. Korgaonkar A, Trivedi U, Rumbaugh KP, Whiteley M (2013) Community surveillance enhances *Pseudomonas aeruginosa* virulence during polymicrobial infection. *Proc Natl Acad Sci USA* 110(3):1059–1064. <https://doi.org/10.1073/pnas.1214550110>
177. Murray JL, Connell JL, Stacy A, Turner KH, Whiteley M (2014) Mechanisms of synergy in polymicrobial infections. *J Microbiol* 52(3):188–199. <https://doi.org/10.1007/s12275-014-4067-3>
178. Siddiqui AR, Bernstein JM (2010) Chronic wound infection: facts and controversies. *Clin Dermatol* 28(5):519–526. <https://doi.org/10.1016/j.clindermatol.2010.03.009>
179. Sanchez CJ Jr, Mende K, Beckius ML, Akers KS, Romano DR, Wenke JC, Murray CK (2013) Biofilm formation by clinical isolates and the implications in chronic infections. *BMC Infect Dis* 13:47. <https://doi.org/10.1186/1471-2334-13-47>
180. Noto MJ, Burns WJ, Beavers WN, Skaar EP (2017) Mechanisms of pyocyanin toxicity and genetic determinants of resistance in *Staphylococcus aureus*. *J Bacteriol* 199(17):e00221–17. <https://doi.org/10.1128/jb.00221-17>
181. DeLeon S, Clinton A, Fowler H, Everett J, Horswill AR, Rumbaugh KP (2014) Synergistic interactions of *Pseudomonas aeruginosa* and *Staphylococcus aureus* in an in vitro wound model. *Infect Immun* 82(11):4718–4728. <https://doi.org/10.1128/IAI.02198-14>
182. Peters BM, Jabra-Rizk MA, O'May GA, Costerton JW, Shirtliff ME (2012) Polymicrobial interactions: impact on pathogenesis and human disease. *Clin Microbiol Rev* 25(1):193–213. <https://doi.org/10.1128/CMR.00013-11>
183. Brileya KA, Camilleri LB, Zane GM, Wall JD, Fields MW (2014) Biofilm growth mode promotes maximum carrying capacity and community stability during product inhibition syntrophy. *Front Microbiol* 5:693. <https://doi.org/10.3389/fmicb.2014.00693>
184. Hillesland KL, Stahl DA (2010) Rapid evolution of stability and productivity at the origin of a microbial mutualism. *Proc Natl Acad Sci USA* 107(5):2124–2129. <https://doi.org/10.1073/pnas.0908456107>
185. de Mazancourt C, Schwartz MW (2010) A resource ratio theory of cooperation. *Ecol Lett* 13(3):349–359. <https://doi.org/10.1111/j.1461-0248.2009.01431.x>
186. Wang M, Schaefer AL, Dandekar AA, Greenberg EP (2015) Quorum sensing and policing of *Pseudomonas aeruginosa* social cheaters. *Proc Natl Acad Sci USA* 112(7):2187–2191. <https://doi.org/10.1073/pnas.1500704112>
187. Allegretta G, Maurer CK, Eberhard J, Maura D, Hartmann RW, Rahme L, Empting M (2017) In-depth profiling of MvfR-regulated small molecules in *Pseudomonas aeruginosa* after quorum sensing inhibitor treatment. *Front Microbiol* 8(MAY):1–12. <https://doi.org/10.3389/fmicb.2017.00924>
188. Filkins LM, Graber JA, Olson DG, Dolben EL, Lynd LR, Bhujar S, O'Toole GA (2015) Coculture of *Staphylococcus aureus* with *Pseudomonas aeruginosa* Drives *S. aureus* towards fermentative metabolism and reduced viability in a cystic fibrosis model. *J Bacteriol* 197(14):2252–2264. <https://doi.org/10.1128/jb.00059-15>
189. Das T, Kutty SK, Tavallaie R, Ibugo AI, Panchompoo J, Sehar S, Aldous L, Yeung AW, Thomas SR, Kumar N, Gooding JJ, Manefield M (2015) Phenazine virulence factor binding to extracellular DNA is important for *Pseudomonas aeruginosa* biofilm formation. *Sci Rep* 5:8398. <https://doi.org/10.1038/srep08398>
190. Glasser NR, Kern SE, Newman DK (2014) Phenazine redox cycling enhances anaerobic survival in *Pseudomonas aeruginosa* by facilitating generation of ATP and a proton-motive force. *Mol Microbiol* 92(2):399–412. <https://doi.org/10.1111/mmi.12566>
191. Deziel E, Lepine F, Milot S, He J, Mindrinos MN, Tompkins RG, Rahme LG (2004) Analysis of *Pseudomonas aeruginosa* 4-hydroxy-2-alkylquinolines (HAQs) reveals a role for 4-hydroxy-2-heptylquinoline in cell-to-cell communication. *Proc Natl Acad Sci USA* 101(5):1339–1344. <https://doi.org/10.1073/pnas.0307694100>
192. Meirelles LA, Newman DK (2018) Both toxic and beneficial effects of pyocyanin contribute to the lifecycle of *Pseudomonas aeruginosa*. *Mol Microbiol* 110(6):995–1010. <https://doi.org/10.1111/mmi.14132>
193. Hall S, McDermott C, Anoopkumar-Dukie S, McFarland AJ, Forbes A, Perkins AV, Davey AK, Chess-Williams R, Kiefel MJ, Arora D, Grant GD (2016) Cellular effects of pyocyanin, a secreted virulence factor of *Pseudomonas aeruginosa*. *Toxins (Basel)* 8(8). <https://doi.org/10.3390/toxins8080236>
194. Price-Whelan A, Dietrich LE, Newman DK (2006) Rethinking 'secondary' metabolism: physiological roles for phenazine antibiotics. *Nat Chem Biol* 2(2):71–78. <https://doi.org/10.1038/nchembio764>
195. Hoffman LR, Deziel E, D'Argenio DA, Lepine F, Emerson J, McNamara S, Gibson RL, Ramsey BW, Miller SI (2006) Selection for *Staphylococcus aureus* small-colony variants due to growth in the presence of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci* 103(52):19890–19895. <https://doi.org/10.1073/pnas.0606756104>
196. Mashburn LM, Jett AM, Akins DR, Whiteley M (2005) *Staphylococcus aureus* serves as an iron source for *Pseudomonas aeruginosa* during in vivo coculture. *J Bacteriol* 187(2):554–566. <https://doi.org/10.1128/JB.187.2.554-566.2005>
197. Kinnersley M, Wenger J, Kroll E, Adams J, Sherlock G, Rosenzweig F (2014) Ex uno plures: clonal reinforcement drives evolution of a simple microbial community. *PLoS Genet* 10(6):e1004430. <https://doi.org/10.1371/journal.pgen.1004430>
198. Kinnersley MA, Holben WE, Rosenzweig F (2009) *E. coli* Unibus plurum: genomic analysis of an experimentally evolved

- polymorphism in *Escherichia coli*. PLoS Genet 5(11):e1000713. <https://doi.org/10.1371/journal.pgen.1000713>
199. Bernstein HC, Paulson SD, Carlson RP (2012) Synthetic *Escherichia coli* consortia engineered for syntrophy demonstrate enhanced biomass productivity. J Biotechnol 157(1):159–166. <https://doi.org/10.1016/j.jbiotec.2011.10.001>
 200. Beck AE, Hunt K, Bernstein HC, Carlson RP (2016) Interpreting and designing microbial communities for bioprocess applications, from components to interactions to emergent properties. In: Eckert CA, Trinh CT (eds) Biotechnology for biofuel production and optimization. Elsevier, Amsterdam, pp 407–432
 201. Carlson RP, Oshota OJ, Taffs RL (2012) Systems analysis of microbial adaptations to simultaneous stresses. In: Wang X, Chen J, Quinn P (eds) Reprogramming microbial metabolic pathways. Subcellular Biochemistry, vol 64. Springer, Dordrecht. https://doi.org/10.1007/978-94-007-5055-5_7
 202. Schepens D, Beck AE, Heys JJ, Gedeon T, Carlson RP (2017) The Benefits of resource partitioning and division of labor in microbial consortia. In: P Amar FK, V Norris (ed) Advances in Systems and Synthetic Biology EDP Sciences Publishing, pp 137–148
 203. Schepens D, Carlson RP, Heys J, Beck AE, Gedeon T (2019) Role of resource allocation and transport in emergence of cross-feeding in microbial consortia. J Theor Biol 467:150–163. <https://doi.org/10.1016/j.jtbi.2019.01.030>
 204. Patel A, Carlson RP, Henson MA (2019) In silico metabolic design of two-strain biofilm systems predicts enhanced biomass production and biochemical synthesis. Biotechnol J 14(7):e1800511. <https://doi.org/10.1002/biot.201800511>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.